

Carbon accumulation potentials of post-SMCRA coal-mined lands¹

C.E. Zipper, J.A. Burger, J.M. McGrath, B. Amichev²

Abstract: Many coal-surface mines reclaimed under SMCRA in eastern US were not restored to forest vegetation and are not currently in managed use. Reforestation of these lands could provide benefits including timber production, watershed protection, and carbon (C) sequestration. Our objectives were to determine the suitability of eastern US coal-mined lands' soil properties for reforestation and to estimate the cumulative potential of these lands to accumulate C through reforestation. Databases of coal mining permits issued under SMCRA were obtained for KY, OH, VA, and WV, and 20 bond-released permits were selected in each state using an area-weighted randomization procedure. Access permissions were obtained for 25 sites (6 each in OH, KY, and VA, and 7 in WV), each of which was sampled at up to 10 randomly selected points. At each sampling point, soil physical properties were determined for the top 30 cm, and soil chemical properties were determined for surface (0 – 10 cm) and subsurface (10 – 30 cm) soil layers. Measured soil properties were used to estimate forest site productivity (SI50) using two methods, and each site's potential to accumulate C was estimated as a function of SI50 based on relationships derived from soil property and C accumulations on pre-SMCRA surface mines. Assuming these sites to be representative of eastern US mined lands, and that 50% of eastern US post-SMCRA mined lands could be available for reforestation under sufficient financial incentives, post-SMCRA mined lands reforested with pine species have the potential to accumulate on the order of 1.6 Tg C yr⁻¹ over a 30 year rotation, equivalent to about 0.2% of projected US coal-combustion C emissions.

Introduction

Large quantities of carbon dioxide (CO₂) and other infrared-absorbing “greenhouse” gases (GHGs) are being emitted to the atmosphere globally by fossil-fuel combustion for energy utilization and other activities. Scientific observations indicate atmospheric GHG concentrations are rising steadily, creating potential influence on climate and consequent negative impacts on the global environment and economy.

Given the dependence of current living standards in industrialized nations on energy usage, growing populations and increasing energy utilization in emerging economies, and the limitations to fossil-fuel alternatives as energy sources available for near-term and widespread implementation, many have called for measures to reduce energy-related CO₂ and other GHG emissions to the atmosphere while offsetting (or sequestering) some portion of those GHG emissions that continue to occur. While the long-term solution to global climate problem is likely to include technologies such as C capture from fossil-fuel combustion and geologic storage, these technologies remain under development. In contrast, sequestration

¹A longer version of this paper was presented at the 2007 National Meeting of the American Society of Mining and Reclamation, and published in the conference proceedings, 30 Years of SMCRA and Beyond, June 2-7, 2007, Gillette WY R.I. Barnhisel (ed) published by American Society of Mining and Reclamation, 3134 Montavesta Rd., Lexington, KY 40502.

²C.E. Zipper, Associate Professor, Department of Crop and Soil Environmental Sciences, Virginia Tech, Blacksburg VA 24061, czip@vt.edu; J.A. Burger, Professor, Department of Forestry, Virginia Tech, Blacksburg VA 24061, jaburger@vt.edu; J.M. McGrath, Assistant Professor, Environmental Science & Technology, 0214 H.J. Patterson Hall, University of Maryland, College Park, MD

methods that rely on management of agricultural and forested systems (terrestrial sequestration) can be implemented with current technologies.

This study assesses the potential for conversion of lands mined under the Surface Mining Control and Reclamation Act (SMCRA) to forest for the purpose of sequestering atmospheric C. Research objectives were to determine the suitability of eastern US coal-mined lands' soil properties for reforestation and to estimate the potential of these lands to accumulate C if reforested.

Experimental Methods

Identify and Sample Mine Sites

Twenty bond-released SMCRA mine permits each in Kentucky, Ohio, Virginia, and West Virginia were selected from permitting databases maintained by state agencies using a randomized procedure with each permit weighted by acreage. An effort was made to identify and contact landowners so as to obtain site access permissions. For sites where access permission could not be obtained but knowledgeable parties communicated opportunities to access nearby sites with similar characteristics, these sites were sampled as substitutes for the original sites. Sampling locations were located on each site using a randomization procedure.

Field parameters measured at each sampling location include depth of penetration by a hand-operated screw auger ("auger depth") (Jones *et al.* 2005). Herbaceous biomass was sampled 1 x 1/4 m quadrats (3 per sampling site) and oven dry weights were obtained. Vegetation characteristics within a 20m radius of the sampling site were estimated as proportionate coverages representing bare ground, open water, herbaceous species, shrub species, and forest species; dominant species, species mixes, and/or species groups within each vegetative component were also recorded; where dominant communities varied within the 20-m radius, the area was sketched and the dominant species, mixes, and/or groups within the each sub-area were recorded. Estimated percent compositions recorded for each sampling location were averaged by mine site.

Soils were sampled by excavation of an approximate 30 x 30 x 30 cm cylindrical volume, separated at 10cm depth into surface and subsurface fractions. The volume of the excavated cavity was determined by filling the hole with small spheres and determining volume occupied by the spheres upon retrieval. Soil samples were bagged and weighed, and a subsample of the soil-sized fraction was removed to determine moisture content. After air drying, soil samples were sieved at 2 mm and the coarse fragment content (>2mm) was determined as a mass proportion of the total sample. Bulk density of the soil-sized fraction was determined as the dry-mass-to-volume ratio, assuming a coarse-fragment density of 2.4 g/cm³. Particle size distribution of the soil-sized fraction (% sand, % silt, % clay of the soil-sized fraction) was determined using the hydrometer method (Gee and Bauder, 1986). Bulk density of the soil-sized fragments was determined on a whole-soil basis (0 – 30 cm), while coarse fragment content and composition were determined separately for 0 -10 and 10 – 30cm depths. Particle size analysis was conducted on a composite sample (1/3 surface + 2/3 subsurface) of soil-sized fragments.

All soil chemical properties were determined separately for the surface and subsurface soil fractions. Soil pH was determined on a mixed suspension of 1:1 volume-to-volume ratio of soil material to distilled water (McLean, 1986). Soluble salts were determined using the method of Rhoades (1986) by measuring the electrical conductivity of a 1:2 volume:volume ratio of soil material to distilled water, and converted to a 1:5 soil:water ratio basis using a regression equation developed by measuring of 1:5 soil:water EC of 32 of the collected samples selected using a randomized procedure biased towards higher values so as to represent the full range of EC measurements.

Sodium bicarbonate extractable P was determined using the method of Olsen and Summers (1986), and exchangeable cations (Ca, Mg, K, Na) were extracted with 1 M ammonium acetate (Thomas, 1986). Phosphorus and cations were determined using a Thermo Elemental ICAP 61E (Inductively Coupled Argon Plasma Atomic Emission Simultaneous Spectrometer) by Virginia Tech Soil Testing Laboratory.

Exchangeable acidity was determined using the potassium chloride extraction technique (Thomas, 1986). Effective cation exchange capacity (ECEC) was estimated by summing the charge associated with exchangeable acidity and exchangeable Ca, Mg, K, and Na. Base saturation was calculated as the proportion of the ECEC occupied by base cations (Thomas, 1986). Total C and N were determined using a Leco C / N analyzer (LECO Corp, St. Joseph, MI).

The forest productivity of each sampling site was estimated as 50-year site index for white oak (expressed in meters) using two methods.

Estimate Forest Productivities: Jones Model

Forest productivity “sufficiency curves” developed by Jones *et al.* (2005) were used to determine sufficiency levels for each soil measurement at each sampling location and a forest site productivity index was determined for each sampling location, also using Jones *et al.* (2005). Sufficiency values were assigned to soil parameters at each sampling location on a 0 – 1 scale, with 0.0 indicating poor forest site quality and 1.0 indicating superior quality. These values were combined to calculate a productivity index using the following equation:

$$PI_{JONES} = 0.44 * S_{BD} + 0.28 * S_{AD} + 0.20 S_{s+c} + 0.08 S_{PH} \quad (1)$$

Where:

- PI_{JONES} = Productivity Index (Jones method).
- S_{BD} = Bulk density of the soil-sized fraction (g/cm³) sufficiency.
- S_{AD} = Augur depth (cm) sufficiency.
- S_{s+c} = Soil texture (% of soil-sized fraction comprised of silt + clay) sufficiency.
- S_{PH} = Soil pH sufficiency.

Jones et al (2005) developed and validated this method using samples taken from the top 20 cm. Our sampling depths were 30 cm. For properties analyzed for the full 30 cm depth (texture, bulk density), measured values were used to determine sufficiency levels. For properties analyzed separately for 0-10 cm and 10-30 cm layers (pH), a “whole soil” value was calculated by considering component values as proportionate to sampling depth.

A mean PI value was calculated for each mine site, and state and overall means were calculated from the mine-site means. The productivity index for each sampling site was converted to a 50-year site index for white pine, expressed in feet (SI50_{WP}), using an equation derived by Jones *et al.* (2005), Figure 7:

$$SI50_{WP} = 117.74 * PI_{JONES} + 15.588 \quad (2)$$

An equivalent 50-year white oak site index expressed in feet (SI50_{WO}) was calculated for each sampling location, using a relationship derived from Figure 1 in Doolittle (1958):

$$SI50_{WO} = 0.8627 * SI50_{WP} - 5.7647 \quad (3)$$

Estimate Forest Productivities

Because the Jones method for estimating forest productivity relies primarily on physical parameters, utilizes a measure that proved difficult to apply with consistent results in the field (auger depth), and was validated using measurements from only a single species (eastern White Pine), the investigators developed an alternative method for estimating soil productivity, termed hereafter as the General Model.

Based on research experience (including Andrews et al, 1992; Jones *et al.* 2005; Showalter *et al.* 2006; Torbert *et al.* 1988; Torbert *et al.* 1992; and other studies), an expression for calculating a productivity index (PI_{GEN}) was developed (see Table 1):

$$PI_{GEN} = (S_{BD} + S_{CF} + S_{TX} + S_{BS} + S_{pH} + S_{EC} + S_p) / 7 \quad (4)$$

Equations used to generate sufficiency values are listed in Table 1. The adequacy of the PI_{GEN} as an indicator of site index was validated using data collected by Rodrigue (2001; see also Rodrigue and Burger 2004) in his study of forest site productivity in mined land forest sites in eastern USA. Using Rodrigue's data (mined sites only, excluding non-mined control sites; 3 to 4 sampling locations per site, at 11 sites in IL, KY, OH, PA, and WV, with the VA site excluded from analysis because data were incomplete and the IN sites excluded because of uneven aged stand characteristics), sufficiency values and a PI_{GEN} value were calculated for each sampling location, and a mean PI_{GEN} was calculated for each site. Rodrigue had measured site index (SI) for the dominant species at each site, and converted that measured SI value to a 50-year white-oak site index (SI_{50WO}). PI_{GEN} values were regressed against SI_{50WO} to test the validity of PI_{GEN} as a site-index indicator, and to define an SI_{50} predictive relationship.

The forest productivity of each of the current study's sampling sites was estimated by applying the General Model method to whole-soil values. For properties analyzed over the full 30 cm sampling depth (texture, bulk density), measured values were used as estimates of whole-soil values and to determine sufficiency levels. For properties analyzed separately for 0-10 cm 10-30 cm layers (pH), a whole-soil value was calculated assuming a depth of 78 cm (as per Andrews *et al.* 1992); the 0-10 cm value was considered as representative of 0–10 cm layer, the 10-30 cm value was considered as representative of soil properties extending from 10 – 78 cm, and a whole-soil value was calculated by considering component values as proportionate to representative depth (*i.e.*, with the 0-10 cm measurement contributing 12.8% to the whole-soil value). Sufficiency levels were determined using the whole-soil values.

Table 1. Sufficiency variables, and calculation methods, used to estimate a soil productivity index (PI_{GEN}) in the general model.

Sufficiency			
Variable	Soil Measure	Units	Sufficiency Calculation
S_{BD}	Bulk Density	$g\ cm^{-3}$ (soil sized fragments)	If $BD < 1.20$, $S_{BD} = 1.0$ If $BD\ 1.20-1.39$, $S_{BD} = 2.20 - 1.0 * BD$ If $BD\ 1.40-1.70$, $S_{BD} = 4.53 - 2.67 * BD$
S_P	Extractable P	mg/kg	If $P > 9$, $S_P = 1.0$ If $P < 9$, $S_P = 0.29 + 0.32 * (\ln P)$
S_{EC}	Electrical conductivity	$dS\ m^{-1}$ (1:5 soil:water ratio)	If $EC < 0.05$, $S_{EC} = 1.0$ If $EC > 0.05$, $S_{EC} = 1.04211 - 0.84210 * EC$
S_{PH}	pH	Standard units	If $pH < 5$, $S_{PH} = (-1) + 0.4 * pH$ If $pH > 5$ and < 6.5 , $S_{PH} = 1.0$ If $pH > 6.0$, $S_{PH} = 3.17 - 0.33 * pH$
S_{CF}	Coarse Fragments	% of soil mass	If $CF < 30$, $S_{CF} = 0.6 + 0.0133333 * CF$ If $CF > 30$ and $< 50\%$, $S_{CF} = 1.0$ If $CF > 50$, $S_{CF} = 2.666667 - 0.03333 * CF$
S_{TX}	Silt + Clay	% by mass of <2mm fragments	If $(s+c) < 35$, $S_{TX} = 0.1 + 0.025791 * (s+c)$ If $s+c > 35$ and $< 60\%$, $S_{TX} = 1.0$ If $(s+c) > 60$, $S_{TX} = 1.9 - 0.015 * (s+c)$
S_{BS}	Base Saturation	%	If $BS < 40$, $S_{BS} = 0.2 + 0.02 * BS$ If $BS > 40$, $S_{BS} = 1.0$

Estimate C Accumulation Potentials of Sampled Mine Sites:

C accumulation potentials for each mine site were estimated using equations derived by Amichev (2007) using data collected by Rodrigue (2001) as described by Burger *et al.* (2006):

For pines: $Eco_C = \text{Exp}(4.416+0.779*SI/\text{age})$ (5)

For mixed stands: $Eco_C = \text{Exp}(4.147+1.039*SI/\text{age})$ (6)

For hardwoods: $Eco_C = 219.455-3711.777*(1/\text{age})$ (7)

Eco_C = C accumulation as above- and below-ground biomass, litter, and soil organic materials on reforested mine sites (Mg C ha^{-1}).

SI = 50 year white-oak site index equivalent, expressed in meters (Because a number of different species were found growing on the pre-SMCRA sites, Amichev standardized all SI's to a 50-year white-oak equivalent, using Doolittle (1958)).

Age = Average age, in years, of the dominant and co-dominant trees.

Estimate C Accumulation Potentials and Significance of Post-SMCRA Mine Lands:

The total area of land disturbed and reclaimed by mining operations in eastern US states under SMCRA was estimated from SMCRA bond-release data obtained from US OSMRE. The distribution of soil and site conditions, and forest productivities, on the sampled mine sites were considered to be representative of the distribution of soil and site conditions on post-SMCRA mined areas. Considering these data, information gained through consultation with state and federal mining agency officials, the investigators' experience, and best professional judgment, the proportion of reclaimed mined areas potentially "available" for reforestation was estimated and applied to the SMCRA bond-release acreage totals to estimate the total acreage of post-SMCRA mined and reclaimed lands potentially "available" for reforestation. The total amount of C that could potentially be accumulated on these lands if reforested was estimated by applying equations (5), (6), and (7). The potential significance of these lands' C accumulation potentials was estimated assuming lands would be reforested over a multiple-year period centered on the year 2010.




Figure 1. Mine site sampling locations. Numbers correspond with mine-site numbers on Table 2.

Table 2. Mine sites where soils and vegetation were sampled and characterized.

Mine Site	Number of Samples	County	Reclam.. Date [†]	Predominant Cover / Land Use
KY-1	5	Breathitt	1990	Abandoned grasses
KY-2	10	Wolfe	1996	Abandoned grasses
KY-3	10	Magoffin	1996	Abandoned grasses
KY-4	10	Knox	1990	Mixed grasses & scrub woody (black locust)
KY-5	3	Breathitt	1996	Reforested with pines
KY-6	10	Perry	1995	Abandoned grasses
OH-1	10	Muskingum	1987	Abandoned grasses
OH-2	9	Vinton	1990	Abandoned grasses
OH-3	10	Coshocton	2000	Abandoned grasses
OH-4	10	Gallia	1985	Wildlife Habitat
OH-5	9	Muskingum	1986	Wildlife Habitat
OH-6	6	Tuscarawas	1980	Grasses (Pastured until recently)
VA-1	8	Buchanan	1989	Grasses & scrub woody (black locust, red maple)
VA-2	9	Buchanan	1986	Forest - mixed hardwood & pines
VA-3	10	Buchanan	1988	Forest - mixed hardwood & pines
VA-4	10	Russell	1998	Mixed white pine & scrub woody (black locust)
VA-5	10	Dickenson	1995	Grasses (Pastured until recently)
VA-6	10	Wise	1998	Mixed grasses & scrub woody (autumn olive)
WV-1	10	Preston	1990	Abandoned grassland / scrub woody
WV-2	9	Mingo	1985	Scrub woody (mixture of planted and invasive hardwoods, black locust, and autumn olive).
WV-3	10	Logan	1994	Grasses & scrub woody (b. locust, autumn olive)
WV-4	8	Logan	1987	Mixed grasses & black locust.
WV-5	9	Logan	1995	Abandoned grasses
WV-6	10	McDowell	1984	Grasses & scrub woody (autumn olive & other)
WV-7	10	Wyoming	1990	Scrub woody (black locust and other)

[†] Estimated average for sampling locations, based on available information as explained in text.

Results and Discussion

Six mine sites were sampled in each of three states (OH, KY, and VA) and seven mines were sampled in WV (Figure 1). The 25 sites yielded 225 sample locations (Table 2).

Vegetation

Vegetative cover characteristics were highly variable (Table 3). In order to compute mine site means, dominant vegetation species and communities recorded were grouped. The predominant planted-herbaceous species on all sites was tall fescue (*Schedonorus phoenix* (Scop.) Holub). Although sericea lespedeza (*Lespedeza cuneata* (Dum.-Cours.) G. Don) was present as a minor component of the vegetation mix on many sites, especially on VA, KY, and WV, it was rarely dominant or co-dominant. The highest percentage cover for planted herbaceous species was computed for OH, but no statistical differences occurred between states. Numerous species – including choke berry (*Photinia sp.*), greenbrier (*Smilax sp.*), poison ivy (*Toxicodendron radicans* L.), ladies lace (*Daucus carota* L.), sumac (*Rhus sp.*), and Virginia creeper (*Parthenocissus quinquefolia* L.) -- were found on single or small numbers of sites

Table 3. Mine-site mean values of estimated percent cover provided by dominant vegetation types at sampling locations among mine sites, by state and for all mine sites sampled.

State - Site	- Planted Herbaceous -				- - - Native Woody - - -								
	Fes- cue	Oth- er gras ses	Lesp- e- deza	Other herba- ceous Leg- umes	Na- tive herb, shrub, vine	Black locust	Red maple	Other native hard- wood	Native soft- wood	Inva- sive	Oth- er	Ba- re	Water
KY	36	2	12	16	4	10	0	3	8	2	-	7	-
OH	50	8	5	12	20	0	-	2	-	2	0	-	0
VA	20	-	13	3	9	7	10	7	17	12	0	2	-
WV	34	0	8	5	10	20	4	3	2	10	-	3	-
All	35	2	9	9	11	10	4	4	7	7	0	3	0

Table 4. Numbers of samples collected, mean values for herbaceous cover and soil physical properties at sampling locations by mine site, by state[†].

State- Site	Sam- ples	Herbaceous cover (kg ha ⁻¹)	Auger depth (cm)	Bulk Density (g cm ⁻³)	Coarse Fragments (%)	Silt + Clay (% of soil-size frags)
KY	48	1217 ^a	27 ^a	1.13 ^a	56 ^a	62 ^{ab}
OH	54	1010 ^a	33 ^a	1.24 ^a	38 ^b	73 ^a
VA	57	1143 ^a	34 ^a	1.17 ^a	51 ^a	59 ^{ab}
WV	66	1140 ^a	25 ^a	1.26 ^a	54 ^a	53 ^b
All	225	1136	30	1.21	50	62

[†] State means followed by different letters are significantly different ($\alpha = .05$), as determined via Tukey's HSD test applied to site means.

and included in the “native herbaceous, shrubs, and vines” category, which represents non-forest-tree native species that were unlikely to have been planted and apparently invaded from surrounding terrain. Blackberry (*Rubus sp.*) was identified as a dominant or co-dominant shrub species at one or more sampling locations on 13 of the 25 sites. Golden rod (*Solidago sp.*), ragweed (*Ambrosia sp.*), thistle (*Cirsium sp.*), and aster (*Aster sp.*) were the primary species recorded as dominant on the Ohio sites.

Few woody species were found on the OH mine sites. On the VA, WV, and KY mine sites, black locust was the most common native woody species found, which is not surprising since it is widely planted. Red maple was recorded commonly in WV and VA in patterns that appear to have occurred through natural invasion. Other native woody species with the potentials to form harvestable tree crops – including Virginia pine, eastern white pine, and hardwood species such as oaks and tulip poplar – were found, with the greatest frequencies in VA and WV. Native woody species were recorded as occurring more commonly as dominant and co-dominant vegetative components at VA sites than at OH sites.

Autumn olive was the most common invasive species recorded, followed by muliflora rose.. As a non-native species that is easily spread by birds, Autumn olive is classified as an “invasive” (Table 3) although some of its occurrences are likely to have originated as reclamation plantings. Autumn olive was recorded as a dominant or co-dominant component on all VA and WV sites, often at several locations. On sites where herbaceous biomass was sampled, amounts ranged from 506 to 1784 kg ha⁻¹, and the mean was 1136 kg ha⁻¹ (Table 4). No statistical differences among states were noted. Herbaceous biomass was not collected at two Virginia sites where forest cover had closed canopy.

Table 5. Mean (all but C and N) and median (C and N) values for soil chemical properties[†] at sampling locations by mine site, and mean values of mine site means by state and total.[‡]

State-Site	pH	EC (dS m ⁻¹)	CEC (meq /100 g)	BS (%)	P (mg kg ⁻¹)	C (Mg ha ⁻¹)	N (Mg ha ⁻¹)
KY	5.58 ^a	0.17 ^a	8.8 ^a	88 ^a	2.51 ^a	24 ^a	1.0 ^b
OH	6.49 ^a	0.05 ^a	15.1 ^a	96 ^a	4.28 ^a	29 ^a	1.8 ^a
VA	5.73 ^a	0.05 ^a	7.8 ^a	83 ^a	3.59 ^a	20 ^a	1.1 ^b
WV	5.58 ^a	0.03 ^a	6.8 ^a	86 ^a	3.22 ^a	28 ^a	1.2 ^b
All Sites	5.92	0.07	9.7	88	3.35	25	1.2

[†] Soil properties expressed for top 30 cm for C and N, and on a whole soil basis, assuming 78 cm depth for other properties listed.

[‡] State means followed by different letters are significantly different ($\alpha = .05$), as determined via Tukey's HSD test applied to site means for normally distributed variables and to site ranks for non-normally distributed variables (EC, CEC, and BS).

Table 6. Mine site mean values of sufficiency values (S), the productivity index used to estimate site index using the Jones model (PI_{JONES}), and estimated 50 year site-index values calculated using the Jones Model (SI_{50WP}) with mine site means averaged by state and for all mine sites sampled.

State	S _{BD}	S _{AD}	S _{S+C}	S _{pH}	PI _{JONES}	SI _{50WO}
KY	0.89	0.62	0.66	0.75	0.758	25.8
OH	0.83	0.74	0.51	0.70	0.728	24.9
VA	0.84	0.69	0.70	0.86	0.772	26.2
WV	0.78	0.60	0.78	0.88	0.738	25.2
All Sites	0.83	0.66	0.67	0.80	0.749	25.5

Soil Properties:

Although soil properties varied widely among sampling sites, few significant differences were found (Tables 4 and 5). Auger penetration depth proved to be a difficult procedure to implement in a manner that yielded consistent results in the field. Site mean bulk densities of the soil-sized particles ranged from 0.93 to 1.54 g cm⁻³, with a mean value for all sites of 1.21 g cm⁻³, while site-mean coarse fragment contents ranged from 23 to 62% with a mean value of 50%. Silt + Clay (S+C) contents were nominally greatest in OH, an apparent result of that state's requirement that topsoils or soil-like topsoil substitute materials be used to construct a surface medium. Site mean pH values were generally in the range of 5 to 7, with only 4 sites (2 in KY, 1 each in VA and WV) having mean pHs below 5 and only two sites (both in OH) having mean pHs > 7 (Table 5). Measured mean EC values were generally low, relative to ECs typically found in fresh mine spoils, a likely result of the weathering that has occurred in these mine soils. The highest EC values occurred in association with low pH's (< 4.5), indicating a likelihood that continued pyrite weathering is the source of the salts and high EC. Base saturation was generally high at the sampling sites, with 15 of the 25 mine site means at greater than 90%, a likely result of base cations released from the formerly fresh mine soils via soil weathering.

Extractable soil P was highly variable, with mine site means ranging from 1.5 to 6.4 mg/kg and values were well distributed throughout that range. Extractable P levels for the 10-30 cm depth samples at 18 of the 225 sampling locations were recorded as being less than the analytical detection limit, equivalent to 0.78 mg/kg; these values were set at the detection-limit-equivalent value for the purpose of calculating means. More than 50% of the detection-limited soil P values occurred in association with pH>7.

Estimating Forest Productivities

Jones-model sufficiency values and productivity indices were estimated for each sampling site;

sampling site productivity indices were calculated using equation (2), and averaged and transformed to SI50_{wp} values (50 year site index for white pine) by mine site. Auger depth and S+C were interpreted to be the soil properties most limiting to forest productivity (with all-mine-site mean values of 0.66 and 0.67, respectively), according to the Jones model (Table 6).

Soil properties measured by Rodrigue (2001) on older mine sites were translated into sufficiency values and used to calculate PI_{GEN} values for each sampling site, and sampling site values were averaged to calculate a PI_{GEN} value for each mine site. Rodrigue had also estimated SI50_{wo} (50-year site index for white oak) values for each site, based on measured growth rates of the dominant species.

The relationship between PI_{GEN}, estimated from Rodrigue’s (2001) soil data, and SI50_{wo} was found to be positive and statistically significant (p<.05) using Spearman correlation. An equation for estimating SI50_{wo} as a function of PI_{GEN} was generated using .jmp software (Figure 2):

$$Y = \Theta_1 + (\Theta_2 - \Theta_1) / (1 + \exp(\Theta_3*(x-\Theta_4))) \quad (8)$$

Sufficiency values for each component of the General Model were estimated, summed, and used to calculate PI_{GEN} for each sampling location using equation 6; means for each sufficiency value and PI_{GEN} were calculated for each mine site, and mean mine-site values were calculated for each state and for all sites sampled. Using the General model, Extractable P was found to be the soil property that is most limiting to forest productivity (mean Sp = 0.63), while EC and BS were found to be least limiting, with mean sufficiencies of 0.97 and 0.99 respectively. Soil P and BS are indicators of soil nutrient levels. When considered together, these two factors suggest that lack of P, not the base cations (Ca⁺⁺, Mg⁺⁺, and K⁺), tends to create soil fertility limitations to forest growth over time. Mean sufficiencies for the other 4 General Model parameters ranged from 0.80 to 0.88.

Mine-site mean SI50_{wo} values estimated by applying the General Model’s calibration equation (Figure 3) to calculated PI_{GEN} values from ranged from 22.1 to 26.2 m (Table 6), while Jones-model estimated SI50_{wo} values range from 19.9 to 30.0 m (Table 7). Mean values for SI50_{wo} estimated using the two models, however, are quite similar (25.1 m for the General Model, 25.5 m for the Jones Model). Productivity indices and the SI50WOs generated (Figure 4) generated by the two models exhibit highly-significant statistical relationships (p<.001) with one another.




Figure 2. Data used for validation of the General Model.




Figure 3. Relationship of the 50-year site indices (SI50s expressed in meters, below) for mine sites estimated by the Jones Model and the General

Table 7. Mine site mean values of sufficiency values (S) and the productivity index used to estimate site index using the general model (PI_{GEN}), with mine site means averaged by state and for all mine sites sampled.

State	S_{BD}	S_{CF}	S_{S+C}	S_{pH}	S_{EC}	S_{BS}	S_p	PI_{GEN}	$SI50_{WO}$
KY	0.89	0.72	0.89	0.80	0.91	1.00	0.54	0.82	24.4
OH	0.83	0.92	0.79	0.84	0.99	1.00	0.72	0.87	25.2
VA	0.84	0.77	0.91	0.91	0.98	0.97	0.65	0.86	25.2
WV	0.78	0.78	0.92	0.93	1.00	0.98	0.62	0.86	25.6
All	0.83	0.80	0.88	0.87	0.97	0.99	0.63	0.85	25.1

Estimating Cumulative Ecosystem C Potentials of the Mined Land Resource Base:

Several assumptions were made in order to estimate the total C-accumulation potentials of the post-SMCRA mined land resource base:

Assumption 1: The forest productivities of mine sites sampled and characterized are representative of those which occur across eastern US. The four states where sampling sites were located accounted for 78% of the combined Phase III and recent (2001-2005) Phase I bond-released lands in eastern US. The fact that few statistically-significant differences among states in soil properties, and no statistically significant differences among states in site index, were found provide no reason to stratify the analysis by state.

Assumption 2: The quantity of Eastern US mined lands that could be available for reforestation and sequestration is on the order of 300,000 ha. This estimate was derived assuming approximately 1/2 of lands mined and reclaimed under SMCRA could be made available for C sequestration. The 300,000 ha figure is slightly less than half of the total Phase 3 and recent Phase I (2001-2005) bond released acreage tallied by US Office of Surface Mining. These acreages, however, include some double-counting due to re-mining of lands that were mined previously under SMCRA, but no estimate of such double-counted acreages is available. We are estimating that about 50% of available acreage could be made available under the right incentive structure considering several factors. First of all, some of the post-SMCRA mined lands are already in beneficial uses, such livestock grazing, and various forms of development. We are aware of no data to allow an estimate of such a quantity, but our experience indicates it to be very small, far less than 50%.

Second, some land owners may not wish to make their lands available for C sequestration even if

additional incentives were offered. If mineral owners believe that future marketplace changes will make it profitable to mine such properties again, so as to extract coals that could not be exploited economically when past mining was occurring, most would consider potential returns from future mining as likely to far outweigh potential returns from reforestation and C sequestration. We would not expect a majority of landowners to fall into this category.

A third factor considered concerns the status of the vegetative communities on mine sites at present. Assuming soil properties are equally favorable, reforestation will be most feasible where mine sites are not occupied by invasive woody vegetation (such as autumn olive) or by low-value species such as black locust. If such species are present and have advanced to the point where they would hinder reforestation, they must be eliminated if the site is to be successfully reforested; this can only occur at some cost. Although such species do accumulate C, and can accumulate it rapidly during the first 10-20 years after reclamation, those accumulations are generally not harvested and converted to sequestrable forms such as building products.

Assumption 3: Post-reforestation C accumulation on planted sites can be estimated using equations 5, 6, and 7. Carbon accumulations on post-SMCRA mined lands were estimated assuming rotation lengths of 30 years for pines and mixed pine-hardwood stands, and 60 years for hardwoods, as per Amichev (2007).

Carbon Accumulation on Post-SMCRA Mined Lands:

Application of equations 5, 6, and 7 using the above assumptions allows calculation of estimated average C accumulations over single rotations of 160, 152, and 158 Mg C ha⁻¹ for pines, mixed pine-hardwood, and hardwoods, respectively (Table 8). These total accumulations correspond with C accumulation rates of 5.3, 5.1, and 2.6 Mg ha⁻¹ yr⁻¹, respectively, for the three stand types (Table 9).

The assumption that 300,000 ha could be reforested with an appropriate incentive structure over a relatively short time period allows extension of the above to estimate that the cumulative potential of the eastern US mined-land resource base to accumulate C if reforested is less than 2 Tg yr⁻¹, on average over an initial rotation. Assuming that pines and/or mixed stands would be targeted by any such incentive program because of their potential to accumulate C more rapidly than hardwoods, post-SMCRA mined lands cumulative potential to accumulate C is equivalent to about 0.2% of projected US combustion emissions from coal in 2025, the midpoint of a representative 30-year rotation beginning circa-2010 (Table 9). Coal consumption within the eastern mining states that were the subject of this research constituted about 23% of US coal consumption in 2005 (US EIA 2006a). Thus, the estimated potential of mined lands, if reforested with pines, to accumulate C is on the order of 1% of total projected coal-related emissions from these states.

The potential of reforested mine sites to accumulate C, as represented by these findings, should be considered as less than their potential to *sequester* C so as to offset energy-related C emissions. Two factors that have not been considered in generating these C-accumulation estimates are the *additionality* of potential reforested-mine-site C accumulations (*i.e.*, the amount of accumulated C over and above that which would occur in the absence of active reforestation), and proportion of accumulated C that could be converted to climate-neutral forms, such as long-term soil C, wood-products placed in long-term non-biodegradable uses, and renewable fuels used to offset fossil-fuel utilization.

Table 8. Mean estimated C accumulation (Mg C ha⁻¹) over a single rotation for three stand types with average site indices estimated using the general model and the Jones model.

Stand Type	Rotation (years)	Jones	General	Average
Pines	30	159	161	160
Mixed Pines /Hardwoods	30	151	154	152
Hardwoods	60	158	158	158

Table 9. Mean estimated C accumulation for three stand types on sampled mine sites over a single rotation and assuming sampled sites are representative of 300,000 ha with potential for reforestation.

Stand Type	Pines	Mixed	Hardwoods
<i>On Sampled Mine Sites:</i>			
Over 1 Rotation (Mg C ha ⁻¹)	160	152	158
Rotation Length (yr)	30	30	60
Annual Rate (Mg C ha ⁻¹ yr ⁻¹)	5.3	5.1	2.6
<i>Cumulative on 300,000 ha:</i>			
Average Rates (Tg yr ⁻¹)	1.60	1.52	0.79
As proportions of US coal combustion C emissions ^a :			
2004 emissions	0.28%	0.27%	0.14%
2010 projected emissions	0.25%	0.24%	0.12%
2025 projected emissions	0.20%	0.19%	0.10%

^a Emissions data from US EIA 2006b, Reference Case Table A-18.

Summary and Conclusions

Eastern US lands that have been mined for coal and reclaimed under SMCRA constitute a significant land resource. Large proportions of eastern US coal-mined lands have not been reforested and are currently not under active management. We investigated the potential for these lands to accumulate C if actively reforested. Should such reforestation occur, results would also include ancillary benefits as environmental services, such as enhanced watershed protection, and timber products.

Soil properties on these mine sites appear to be generally well-suited for reforestation. The mean 50-year site index for white oak on 25 mine sites identified through a random selection procedure was estimated at about 25 meters using two different methods, a level that would be considered as better-than-average on natural soils occurring in most eastern US areas mined for coal. Soil compaction was not found to be a major limiting factor on most sites, although some sites were heavily compacted. Low amounts of plant-available phosphorous, a soil property that would be easily ameliorated through fertilization, was the property found to be most limiting to reforestation potentials.

One major site factor that occurs commonly and that can be expected to hinder reforestation of sites where it occurs is the proliferation of low-value woody species with little potential to produce marketable timber – especially black locust and autumn olive. When these species reach advanced stages, the cost of removal can render reforestation investments unprofitable. As time passes, it is likely that many mine sites which are not currently occupied by such species in advanced growth stages will become so occupied and such sites' potential for reforestation will decline as a direct result.

If on the order of 50% of the lands mined for coal and reclaimed under SMCRA in the eastern US were to be reforested, estimated rates of C accumulation on such sites are in the range of 0.8 – 1.6 Tg C yr⁻¹, on the order of 0.2% of US C emissions from coal combustion.

Although potential C accumulation and sequestration quantities are not great relative to potential national needs should energy-related C-emissions offsets become a requirement at some future date, these lands are available and unused, and many possess soil and site properties that are well-suited to reforestation. As time passes, their collective potential for reforestation can be expected to decline due to continued proliferation and growth of low-value woody species such as black locust and autumn olive.

Acknowledgements

The authors acknowledge and appreciate the efforts of numerous individuals representing mining firms, landowners, and agency personnel who provided permissions and assistance to research, allowing us to locate and access the sampling sites. Thanks also to Sarah Hash, W.T. Price, Julie Burger, and

Clinton Steele for conducting and assisting laboratory analyses; to Patricia Donovan for analysis of Landsat imagery to estimate southern WV and VA site ages; and Golde Holtzman for his assistance in developing the General Model SI_{50} estimation function. Primary funding support for this work provided by US Department of Energy (DE-FG26-02NT41619), with additional support provided by National Aeronautics and Space Administration (NASA Grant NNS04AB26G) and Powell River Project.

Literature Cited

- Akala, V. A., and R. Lal. 2001. Soil organic carbon pools and sequestration rates in reclaimed minesoils in Ohio. *J. Environ. Qual.* 30:2098–2104.
- Amichev, B. 2007. Biogeochemistry of Carbon on Disturbed Forest Landscapes. Ph.D. dissertation. Department of Forestry, Virginia Tech. In preparation.
- Andrews, J.A., J.E. Johnson, J.L. Torbert, J.A. Burger, and D.L. Kelting. 1998. Minesoil properties associated with early height growth of eastern white pine. *Journal of Environmental Quality* 27:192-198.
- Burger, J.A., J. Galbraith, T. Fox, G. Amacher, J. Sullivan, and C. Zipper. 2006. Restoring Sustainable Forests on Appalachian Mined Lands for Wood Products, Renewable Energy, Carbon Sequestration, and Other Ecosystem Services. Final Report to U.S. Department of Energy, NETL, Instrument No: DE-FG26-02NT41619 (in preparation).
- Doolittle, W. T. 1958. Site index comparisons for several forest species in the southern Appalachians. *Soil Science Society of America Proceedings* 22: 455-458.
- Gee G., J. Bauder. 1986. Particle-size analysis. p. 383-411, in A.K. Klute (ed.) *Methods of soil analysis, part 1*. Soil Sci. Soc. America. Madison, WI
- Jones, A.T. 2005. Site Quality Classification for Mapping Forest Productivity Potential on Mine Soils in the Appalachian Coalfield Region. M.S. Thesis. Virginia Tech. 305 p.
- Jones, A.T., J.M. Galbraith, and J.A. Burger. 2005. A forest site quality classification model for mapping reforestation potential of mine soils in the Appalachian coalfield region. P. 523-539, in: *Proceedings, 2005 National Meeting of the American Society of Mining and Reclamation*. ASMR, 3134 Montavesta Rd., Lexington, KY 40502.
- McLean, E.O. 1986. Soil pH and lime requirement. p. 199–224. in A.K. Klute (ed.) *Methods of soil analysis, part 2*. Soil Sci. Soc. America. Madison, WI
- Nelson D., L. Sommers. 1986. Total carbon, organic carbon, and organic matter. p. 539-589, in A.K. Klute (ed.) *Methods of soil analysis, part 2*. Soil Sci. Soc. America. Madison, WI .
- Olsen, S.R, and L.E. Summers, 1986. Phosphorous. P. 403-430, in A.K. Klute (ed.) *Methods of soil analysis, part 2*. Soil Sci. Soc. America. Madison, WI
- Rhoades, J.D. 1982. Soluble constituents. p. 167–179. In A.K. Klute (ed.) *Methods of soil analysis, part 2*. Soil Sci. Soc. America. Madison, WI
- Rodrigue, J. A. 2001. Woody species diversity, forest and site productivity, stumpage value, and carbon sequestration of forests on mined lands reclaimed prior to the passage of the Surface Mining Control and Reclamation Act of 1977. M.S. Thesis, Virginia Polytechnic Inst. and State Univ., Blacksburg.
- Rodrigue, J. A., and J. A. Burger. 2004. Forest soil productivity of mined land in the Midwestern and eastern coalfield regions. *Soil Sci. Soc. Am. J.* 68(3): 833-844.
- Showalter, J., J. Burger, C. Zipper, J. Galbraith, P. Donovan. 2006. Physical, chemical, and biological mine soil properties influence white oak seedling growth: A proposed mine soil classification model. *Northern Journal of Applied Forestry* (in press).
- Shresthra, R, R. Lal. 2006. Ecosystem carbon budgeting and soil carbon sequestration in reclaimed mine soil. *Environment International* 32: 781–796.
- Thomas, G.W. 1986. Exchangeable cations. p. 159-166, in A.K. Klute (ed.) *Methods of soil analysis, part 2*. Soil Sci. Soc. America. Madison, WI
- Torbert, J. L., A. R. Tuladhar, J. A. Burger, and J. C. Bell. 1988. Minesoil property effects on the height of ten-year-old white pine. *J. Environ. Qual.* 17:189-192.
- Torbert, J. L., J. A. Burger, and W. L. Daniels. 1990. Pine growth variation associated with overburden rock type on a reclaimed surface mine in Virginia. *J. Environ. Qual.* 19:88-92.
- US Energy Information Administration (EIA). 2006a. Annual Coal Report 2005.
- US Energy Information Administration (EIA). 2006b. Annual Energy Outlook 2006.
- Ussiri, D., R. Lal. 2005. Carbon sequestration in reclaimed minesoils. *Critical Reviews in Plant Sciences* 24:151–65.